


Management of Nightmare disorders in adult patients with Post Traumatic Stress Disorder (PTSD)

Note: Prazosin and trazodone are IMOC Amber 2 category – initiation by the sleep centre followed by maintenance prescribing in primary care. GPs are not expected to initiate these treatments but may be asked to take on prescribing in line with IMOC recommendations

Agomelatine is IMOC RED category (hospital only) – initiation and ongoing prescribing will be by the sleep service



General principles of patient management in the Sleep Disorder Centre

- Patients will remain under the consultant that initiated treatment and after the initial reviews will have consultant reviews at 3, 6 and 12 months, then annually.
- Follow up consultations with the sleep pharmacist are available when necessary.
- Patients are counselled on initiation, including when to dose titrate, and when to stop treatment if not tolerated/effective. This detail is clearly communicated to GPs in the clinic letter.
- The initial prescription from the specialist team will be a minimum of 3 months' supply. Patient will be stabilised on medication (with baseline investigations undertaken) before transfer to GP
- When treatment is no longer required (e.g. not effective/tolerated), the GP will be informed via a clinic letter, including any detail of treatment tapering if required
- For further advice please contact: gst-tr.thesleeppharmacistgstt@nhs.net

Approval date: April 2025

Review date: April 2027 (or sooner if evidence or practice)

Not to be used for commercial or marketing purposes. Strictly for use within the NHS

South East London Integrated Medicines Optimisation Committee (SEL IMOC). A partnership between NHS organisations in South East London Integrated Care System: NHS South East London (covering the boroughs of Bexley/Bromley/Greenwich/ Lambeth/Lewisham and Southwark) and GSTFT/KCH /SLaM/ Oxleas NHS Foundation Trusts and Lewisham & Greenwich NHS Trust


Management of Nightmare Disorders in Adult Patients



Drug	Starting dose	Titration instructions	GP Monitoring	Additional comments
Prazosin tablets 	0.5mg at bedtime	<p>Generally, the dose may be increased gradually in steps of 0.5mg fortnightly increments up to maximum of 10mg at night if needed and well tolerated.</p> <p>The sleep disorder Centre may use quicker up-titration methods as determined by the severity of symptoms, and response to treatment.</p>	-Annual BP, HR, renal function & LFTs	<ul style="list-style-type: none"> - Blood pressure to be taken at baseline by Sleep Disorders Centre, and following each dose increment until stable -Patients to be advised for first dose effects. - First dose may cause collapse due to hypotensive effect (therefore should be taken on retiring to bed). - More frequent blood pressure monitoring may be required and dose adjusted accordingly if any concurrent antihypertensives, cardiovascular comorbidities, or hepatic/renal impairment. -Prazosin is indicated for hypertension, and benign prostatic hyperplasia -Prazosin is an alpha-1 adrenergic receptor antagonist that reduces CNS sympathetic outflow throughout the brain. -Several CNS phenomena implicated in the pathogenesis of PTSD are regulated by alpha-1 adrenergic receptors including a number of sleep/nightmare phenomena. -If there are changes to symptoms or monitoring parameters the GP should contact the Sleep Disorders Centre for further advice -discontinue if: symptomatic postural hypotension or risk of precipitating syncope
Trazodone capsules 	50mg at bedtime	<p>Generally, the dose may be increased by 50mg every 2-4 weeks up to 150mg at bedtime, if needed and well tolerated.</p> <p>The sleep disorder Centre may use quicker up-titration methods as determined by the severity of symptoms, and response to treatment.</p>	<ul style="list-style-type: none"> -Repeat ECG if dose changes or prescribed new QTc prolonging medication -Annual BP & HR -Annual renal function, FBCs & LFTs 	<ul style="list-style-type: none"> -Baseline ECG by Sleep Disorders Centre if cardiovascular history or taking QTc prolonging medications. -Trazodone is also indicated for depression. -Depression is associated with PTSD-related nightmares. -Trazodone may be used to manage nightmare disorders when prazosin has failed and patient continues to report sleep initiation insomnia despite improved sleep disturbances. - Trazodone may be used in patients with comorbid insomnia with depression without PTSD-related nightmares at doses up to 300mg at bedtime -If there are changes to symptoms or monitoring parameters the GP should contact the Sleep Disorders Centre for further

Management of Nightmare Disorders in Adult Patients



				<p>advice</p> <ul style="list-style-type: none"> -discontinue if: symptoms of liver disorder, or suicidal ideation <p>SEL pathway Comorbid insomnia in adults SEL formulary recommendation</p>
<p>Agomelatine tablets</p> <p> RED</p>	25mg at bedtime	Increase dose to 50mg at bedtime after 2-4 weeks if needed and well tolerated.	<p>GP monitoring not applicable. For Specialist monitoring.</p> <ul style="list-style-type: none"> -Specialist team to perform LFT monitoring at baseline and thereafter -LFTs before treatment and after 3, 6, 12 and 24 weeks of treatment, and then regularly thereafter when clinically indicated* (restart monitoring schedule if dose increased). -Annual BP & HR -Annual renal function 	<ul style="list-style-type: none"> - Agomelatine is also indicated for depression. -Depression is associated with PTSD-related nightmares. - Agomelatine may be used to manage nightmare disorders when prazosin and trazodone have failed and patient continues to report sleep initiation insomnia despite improved sleep disturbances. - Agomelatine may be used in patients with comorbid insomnia with depression without PTSD-related nightmares -discontinue if serum transaminases exceed 3 times the upper limit of reference range or symptoms of liver disorder * when clinically indicated includes (but not limited to) where dose adjustments may be required due to: (i) worsening nightmare disorder and/or insomnia symptoms, (ii) drug interactions, (iii) smoking started or stopped during treatment, or (iv) concurrent use of drugs associated with hepatic injury <p>SEL pathway Comorbid insomnia in adults SEL formulary recommendation</p>

Approval date: April 2025

Review date: April 2027 (or sooner if evidence or

Not to be used for commercial or marketing purposes. Strictly for use within the NHS

Appendix 1 – Non-pharmacological recommendations

Lifestyle optimisation measures

Stimulus control

- Bed/bedroom use only for sleep, sex and getting dressed (not reading, resting, phone, watching TV etc.)
- Get out of bed if not sleeping after 15-20 min (especially if agitated)

Sleep scheduling

- Limit the time in bed to the average time actually spent sleeping (best achieved via sleep diaries).
- Avoiding daytime and evening naps
- Getting into bed only when sleepy tired
- Getting up at the same time every day, including weekends

Naps

- Discouraged because they reduce sleep drive
- If naps cannot be avoided, then try to limit them to 20 minutes (setting an alarm), and ideally not in the 5 hours before the main episode of sleep

Additional

- Avoid caffeine after 12pm midday; switch to decaffeinated drinks
- Avoid alcohol, at the very least in the 2 hours before bed
- Daily exercise, ideally outside to gain exposure to natural day light
- Other sleep hygiene advice via the [Sleep Foundation](#)

Approval date: April 2025

Review date: April 2027 (or sooner if evidence

Not to be used for commercial or marketing purposes. Strictly for use within the NHS

South East London Integrated Medicines Optimisation Committee (SEL IMOC). A partnership between NHS organisations in South East London Integrated Care System: NHS South East London (covering the boroughs of Bexley/Bromley/Greenwich/ Lambeth/Lewisham and Southwark) and GSTFT/KCH /SLaM/ Oxleas NHS Foundation Trusts and Lewisham & Greenwich NHS Trust